

Binding Energies and Scattering Observables in the $^3\text{He}^4\text{He}_2$ Atomic System*

W. Sandhas^a, E. A. Kolganova^{bc}, A. K. Motovilov^b, and Y. K. Ho^c

^aPI Universität Bonn, Endenicher Allee 11-13, D-53115 Bonn, Germany

^bBLTP, JINR, 141980 Dubna, Moscow Region, Russia

^cIAMS, Academia Sinica, PO Box 23-166, Taipei, Taiwan 10764, R. O. C.

The $^3\text{He}^4\text{He}_2$ three-atomic system is studied on the basis of a hard-core version of the Faddeev differential equations. The binding energy of the $^3\text{He}^4\text{He}_2$ trimer, scattering phase shifts, and the scattering length of a ^3He atom off a ^4He dimer are calculated using the LM2M2 and TTY He-He interatomic potentials.

The system of three He atoms is of great interest for various questions as, e.g., the formation of liquid helium drops, superfluidity or Bose-Einstein condensation. The literature on experimental and theoretical studies of the symmetric ^4He three-atomic system is rather large (see, e. g., [1]–[10] and references cited therein). In contrast, the asymmetric $^3\text{He}^4\text{He}_2$ system has found comparatively little attention. To the best of our knowledge the numerical studies of the $^3\text{He}^4\text{He}_2$ trimers were performed only in Refs. [8]–[11]. Except Ref. [11], there are no scattering calculations reported for this system.

In [5], [6] the symmetric ^4He three-atomic system was studied on the basis of a mathematically rigorous hard-core version of the Faddeev differential equations. Using this approach allows one to overcome technical complications due to the strong repulsive part in the atomic interaction. Along the same line we now study the asymmetric $^3\text{He}^4\text{He}_2$ system. We calculate its bound states, the scattering phase shifts of a ^3He atom off a ^4He dimer at ultra-low energies, and the corresponding scattering lengths.

In the present investigations we employ two realistic He-He atomic interactions, the LM2M2 potential by Aziz and Slaman [12] and the TTY potential by Tang, Toennies and Yiu [13]. As in [5], [6] we use the finite-difference approximation of the two-dimensional partial-wave Faddeev equations. We consider the case of zero total angular momentum and take $\hbar^2/m_{^4\text{He}} = 12.12 \text{ K } \text{\AA}^2$ where $m_{^4\text{He}}$ stands for the mass of a ^4He atom. Notice that the ^4He -dimer energy is 1.30348 mK for the LM2M2 potential and 1.30962 mK for the TTY potential [6].

Due to the smaller mass of the ^3He atom ($m_{^3\text{He}}/m_{^4\text{He}}=0.753517$), the $^3\text{He} - ^4\text{He}$ system is unbound. None the less, the $^3\text{He}^4\text{He}_2$ trimer exists, though with a binding energy one order of magnitude smaller than the one of the $^4\text{He}_3$ ground state (see, e.g. [5] and [6]). And,

*This work was supported by the Deutsche Forschungsgemeinschaft (DFG), the Academia Sinica, the NSC of Taiwan (ROC), and the Russian Foundation for Basic Research

Table 1

Absolute value of the ${}^3\text{He}{}^4\text{He}_2$ trimer binding energy (in mK). Grids used: $N_p = 600$, $N_\theta = 605$, the cut-off hyperradius $\rho_{\text{max}} = 200 \text{ \AA}$.

Potential	l_{max}	This work	[8]	[9]	[10]	[11]
LM2M2	0	7.30	10.22			
	2	13.15				
	4	13.84			13.66	14.4
TTY	0	7.25				
	2	13.09				
	4	13.78			14.165	14.1

in contrast to the symmetric case, there is no excited (Efimov-type) state in the asymmetric ${}^3\text{He}{}^4\text{He}_2$ system.




Figure 1. S-wave ${}^3\text{He}$ atom – ${}^4\text{He}$ dimer scattering phase shifts $\delta_0(E)$ in case of the TTY He–He interatomic potential as a function of the c.m. energy. The lower curve corresponds to the case where $l_{\text{max}} = 0$ while for the upper curve $l_{\text{max}} = 2$. The corresponding phase shift curves for the LM2M2 potential are practically the same.

The best dimensions of the grids employed in our investigation were $N_p = 600$ and $N_\theta = 605$ with a cut-off hyperradius $\rho_{\text{max}} = 200 \text{ \AA}$ (see [5] for explanation of the notation). The maximal value of the angular momenta taken into account in the two-body subsystems was $l_{\text{max}} = 4$. Our results for the ${}^3\text{He}{}^4\text{He}_2$ binding energy as well as the results available in the literature, are presented in Table 1. One can see that at this point the different approaches agree fairly well. A small discrepancy in the results is possibly due to limitations of the computer facilities. We have found that most of the contribution to the ${}^3\text{He}{}^4\text{He}_2$ binding energy stems from the $l = 0$ and $l = 1, 2$ partial waves, about 53% and 42%, respectively. The overall contribution from the

$l = 3$ and $l = 4$ components is of the order of 5 %, that is, approximately the same as in the case of the symmetric ^4He trimer [6]. A certain (but rather small) deepening of the $^3\text{He}^4\text{He}_2$ binding energy may be expected when choosing a larger grid.

Our phase shift results obtained for a grid with $N_\rho = 502$, $N_\theta = 500$, and $\rho_{\max} = 460 \text{ \AA}$ are plotted in Fig. 1. Note that exactly the same grid was employed in the $^4\text{He}-^4\text{He}_2$ phase-shift calculations of [6]. Incident energies below and above the breakup threshold, i. e. both for the elastic scattering ($2+1 \rightarrow 2+1$) and breakup ($2+1 \rightarrow 1+1+1$) processes, were considered. Table 2 contains our results for the $^3\text{He}-^4\text{He}_2$ scattering length.

Table 2

Estimations for the ^3He atom – ^4He dimer scattering length (in $\text{\AA}$). Grids used are the same as in Table 1.

$l_{\max}$	TTY		LM2M2	
	This work	[11]	This work	[11]
0	38.8		38.5	
2	22.4		22.2	
4	21.2	19.6	21.0	19.3

Acknowledgements. We are grateful to Prof. V. B. Belyaev and Prof. H. Toki for providing us with the possibility to perform calculations at the supercomputer of the Research Center for Nuclear Physics of Osaka University, Japan.

REFERENCES

1. S. Grebenev, J.P. Toennies, and A.F. Vilesov: Science **279** (1998) 2083.
2. F. Luo, C. F. Giese, and W. R. Gentry: J. Chem. Phys. **104** (1996) 1151.
3. J. Carbonell, C. Gignoux, and S.P. Merkuriev: Few-Body Systems **15** (1993) 15.
4. V. Roudnev and S. Yakovlev: Chem. Phys. Lett. **328** (2000) 97 (arXiv: physics/9910030).
5. E.A. Kolganova, A.K. Motovilov, and S.A. Sofianos: J. Phys. B **31** (1998) 1279 (arXiv: physics/9612012).
6. A.K. Motovilov, W. Sandhas, S.A. Sofianos, and E.A. Kolganova: Eur. Phys. J. D **13** (2001) 33 (arXiv: physics/9910016).
7. P. Barletta and A. Kievsky: Phys. Rev. A **64** (2001) 042514.
8. B.D. Esry, C.D. Lin, and C.H. Greene: Phys. Rev. A **54** (1996) 394.
9. E. Nielsen, D.V. Fedorov, and A.S. Jensen: J. Phys. B **31** (1998) 4085 (arXiv: physics/9806020).
10. D. Bressanini, M. Zavaglia, M. Mella, and G. Morosi: J. Chem. Phys. **112** (2000) 717.
11. V. Roudnev: Talk at the 17th International IUPAP Conference on Few-Body Problems in Physics (Durham, North Carolina, USA, June 5–10, 2003).
12. R.A. Aziz and M.J. Slaman: J. Chem. Phys. **94** (1991) 8047.
13. K.T. Tang, J.P. Toennies, and C.L. Yiu: Phys. Rev. Lett. **74** (1995) 1546.